



Understanding the selective-sensing mechanism of Al³⁺ cation by a chemical sensor based on Schiff-base. A novel theoretical methodology

Manuel A. Treto-Suárez^a, Yoan Hidalgo-Rosa^a, Ximena Zarate^{b,c*}, Dayan Páez-Hernández^{a, d*}.

^aDoctorado en Fisicoquímica Molecular, Universidad Andrés Bello. Ave. República #275, Santiago de Chile, Chile. <u>mtretosuarez@gmail.com</u>.

^bInstituto de Ciencias Químicas Aplicadas, Facultad de Ingeniería, Universidad Autónoma de Chile, Av. Pedro de Valdivia 425, Santiago, Chile. <u>jazminac@gmail.com</u>.

^cMillennium Nuclei on Catalytic Processes towards Sustainable Chemistry (CSC), Chile.

^dCenter of Applied Nanosciences (CANS), Chile

Abstract

A methodology that allows to explain experimental behavior of a turn-on luminescent chemosensor is proposed and verified in 1-[(1H-1,2,4-triazole-3-ylimino)-methyl]-naphthalene-2-ol] (L1), selective to Al^{3+} cations. This sensor increases its emission when interacting with ion upon excitation at 442 nm, which is denoted as the chelation-enhanced fluorescence (CHEF) effect. The PET is responsible for the fluorescence quenching in L1 at 335 nm, in Ni²⁺/L1 at 385 nm and in Zn²⁺/L1 at 378 nm. In the Ni²⁺/L, The LMCT, from the molecular orbital of the ligand to the Ni $3d_{x^2-y^2}$ orbital, can contributed to the quenching of fluorescence. Based on oscillator strength, the highest luminescent intensity the L1 at 401 nm and the $Al^{3+}/L1$ at 494 nm in relation to the others is evidenced. The consideration of the relative energies of the excited states, the calculation the rate and lifetime of the electron transfer deactivation is necessary to get a good description of the sensor.




Figure. Sensing of Al³⁺ cations, due to the CHEF effect that produce in L1 when is excited at 442 nm

Acknowledgements: The authors thank to Ph.D. Program in Molecular Physical Chemistry from University Andrés Bello, subsidy of the DAD-UNAB, FONDECYT 1180017, FONDECYT 1180565. Millennium Science Initiative of the Ministry of Economy, Development and Tourism-Chile grant Nuclei on Catalytic Processes towards Sustainable Chemistry (CSC).

References.

^[1] J. Jia et al. Tetrahedron Letters 54 (2013) 3471–3474.

^[2] Sarma, S. et al. J. Fluoresc. 25, 1537–1542 (2015).